### 1 Overview

Right off the bat we need to discuss the difference between discrete and continuous. A Discrete unit is indivisible, and we count discrete Theorem 5 (The Extended Pigeon-Hole Principle). If N pigeons are

On the flipside, we measure with continuous units. This gives us

On the impace, we measure with continuous units. In signers us fractions and no-acquire roal numbers.

We also have discrete structures which include sets, sequences, networks, matrices, permutations, and real-world data.

These structures are what the class will focus on.

These structures are what the class will focus on.

is a non-negative integer), then we say the cardinality of S is n, i.e.

If x is an element of S, we say  $x \in S$ .

Theorem 2 (Multiplicative Principle of Counting<sup>2</sup>). If task 1 can be expressed as number of ways to do one task and then the other is n1 · n2.

Theorem 3 (Additive Principle of Counting). If task 1 can be don

2 Principles of Counting

Theorem 4 (The Pigeon-Hole Principle). If n pigeons fly into k pigeon k is k into k pigeon. Example 2.2. Same question as before, but suppose we have one holes, and k < n, then some pigeon hole must contain at least 2 pigeons. chair, one vice-chair, and two treasurers.

 $^{1}$ Cardinality is the number of elements in S. Ordinality is for ordering infinities.  $\hookrightarrow 100 \cdot 99 \cdot {98 \choose 2}$   $^{2}$ Product Rule

Discrete Math Notes

If f is a function from a finite set x to a finite set u, and if |x| > |u|, then  $f(x_1) = f(x_2)$  for some  $x_1, x_2 \in x$  such that  $x_1 \neq x$ 

things. This gives us number such as the set of Natural numbers, assigned to K < N pigeon holes, then one of the pigeon holes must  $\mathbb{N} = \{0, 1, 2, 3, 4, \cdots\}$ .

These structures are what the class will focus on o the structures are what the class will focus on o distinct objects in which order matters.

Theorem 1 (Naive Set Theory). A set is an unordered collection of objects. As a constant of the structure of the

$$P(n, r) = \frac{n!}{(n - r)!} = P_r^n$$

Let A and B be set, the Cartesian product of A and B,  $A \times B$ , In general, if there are n objects, with  $n_1$  of type l,  $n_2$  of type l, ... is the set of all ordered pairs (a,b) where  $a \in A$  and  $b \in B$ , i.e. to type r, then there are  $\frac{all}{a \times b \times b}$  to total permutations of the n objects.  $A \times B = \{(a,b) \mid e A \nmid b \in B\}$ .

Theorem 7. A combinations is a sequence of objects where order does not matter. The size of a combination is the number of different elements that compose it.

The number of combinations of size r using n different objects is

$$C(n, r) = {n \choose r} = C_r^n = \frac{n!}{r!(n - r)!} = \frac{P(n, r)}{r!}$$

APPM 3170

## 2.7.1 Master Theorem Corollary

**Theorem 12.** The generating function for the sequence  $\{a_n\}_{n\geq 0}$  is the

$$A(z) = \sum_{n=0}^{\infty} a_n z^n$$

Think of the zs as placeholders. We don't actually care about their

Is the coefficient of the  $z^{nth}$  term in the series A(z).

 $\hookrightarrow$  Note  $\frac{d}{d}\left(\frac{1}{d}\right) = \frac{1}{d}$ 

Therefore the generating function is  $A(z) = \frac{z}{(1-z)^2}$ 

Theorem 13, If A(z) is the generating function for the sea

 $\left|\bigcup_{i=1}^{n} A_{i}\right| = \sum_{i} |A_{i}| - \sum_{i} |A_{i} \cap A_{j}| + \cdots + (-1)^{n+1} \left|\bigcap_{i=1}^{n} a_{i}\right|$ 

$$P(n, r) = \frac{n!}{(n - r)!} = I$$

Suject Bhat

Let f be an increasing function that satisfies  $f(n)=af(\frac{n}{b})+cn^d$  where  $a,c\in\mathbb{Z}^+,b>1\land c,d\in\mathbb{R},c>0,d\geq 0.$  If  $n=b^k,k\in\mathbb{Z}^+$  then

2. f(n) is  $O(n^d \cdot log(n)) \Leftrightarrow a = b^d$ 3. f(n) is  $O(n^{log_b(a)}) \Leftrightarrow a > b^d$ 

### 2.8 Generating Functions

$$A(z) = \sum_{n=0}^{\infty} a_n z^n$$

Notation:

 $[z^n]A(z)$ 

Example 2.8. Show that the generating function for  $a = n, n \ge 0$  is  $a = n, n \ge 0$ 

But  $\frac{d}{dz} = \left(\frac{1-z}{1-z}\right) = \frac{d}{dz} \left(\sum_{n=0}^{\infty} z^n\right) = \sum_{n=0}^{\infty} n - z^{n-1}$ 

 $z \cdot \frac{1}{(1-z)^2} = z \cdot \sum_{n=0}^{\infty} nz^{-1} = \sum_{n=0}^{\infty} nz^n = \sum_{n=0}^{\infty} a_n z^n \rightarrow a_n = n$ 

associated to  $\{a_n\}_{n\geq 0}$  and if B(z) is the generating func-to  $\{b_n\}_{n\geq 0}$ , then

á. APPM 3170

Discrete Math Notes 5.  $f:A\to B$  is injective, (1-1), or one-to-one, if for any  $y\in B$  there is at most one a in A such that f(a)=y. Big O Form | Complexity 6.  $f : A \rightarrow B$  is surjective, or onto if for any  $y \in B$ ,  $\exists a \in A$  such nlog(n)

7. If f is both one-to-one and surjective, then it is called bijective. 8. A set A is said to be countable if there exists a bijection  $f: \mathbb{N} \to$ 

If f: A → B is a bijection, then it is invertible.

10. A set that is not finite nor countable is said to be uncountable.

# 5 Algorithms and Integers

## 5.1 Complexity

Theorem 19. Definitions

while it is, Definitions,  $F:[0,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , we have  $f:[-,\infty) \to \mathbb{R}$ , we have  $f:[-,\infty) \to \mathbb{R}$ , we have  $f:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , we have  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , and  $g:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , and  $g:[-,\infty) \to \mathbb{R}$ , where  $f:[-,\infty) \to \mathbb{R}$  and  $g:[-,\infty) \to \mathbb{R}$ , an

 $|f(x)| \le c |g(x)|$  for all x > k

2. We write  $f = \Theta(g)$ ,  $\gamma$  and g are of the same order "if f = O(g). A greedy algorithm is an algorithm that makes the "best" choice at and if g = O(f). This is equivalent to saying  $g_{\lambda}$ ,  $g_{\lambda}$ ,  $h(0) \in g_{\lambda}$  c and step,  $g_{\lambda} > h(0) \in G(g_{\lambda})$ .

Theorem 20. If  $f_1(x) = O(g_1(x))$  and  $f_2(x) = O(g_2(x))$ , then

2.  $(f_1f_2)(x) = O(g_1(x) - g_2(x))$ 

### Theorem 21. Definitions.

1. Time Complexity of an algorithm relates to the time required to

- give output.

  2. Space Complexity relates to the computer memory required by the
- 3. Worst-Case Complexity is the maximum number the algorithm

APPM 3170

Theorem 6 (Permutations). A permutation is any linear armnaement

$$P(n, r) = \frac{n!}{(n-r)!} = 1$$

$$C(n,r) = \binom{n}{r} = C_r^n = \frac{n!}{r!(n-r)!} = \frac{P(n,r)}{r!}$$

Theorem 3 (Additive Principle of Conting), If last I can be done
in a wage, and task 2 can be done in a wage, then the other is n<sub>1</sub> + n<sub>2</sub>.

Example 2.1. How many different committees can be formed
consisting of one chair, one vice-chair, and one treasurer from a pool
of 100 people.

2.1 Pigeon-Hole Principle

3.1 Pigeon-Hole Principle

3.2 Pigeon-Hole Principle

3.3 Pigeon-Hole Principle

3.4 How many different committees can be formed
consisting of one chair, one vice-chair, and one treasurer from a pool
of 100 people.

3.4 How many different committees can be formed
consisting of one chair, one vice-chair, and one treasurer from a pool
of 100 people.

3.4 Pigeon-Hole Principle

Ó.

1.  $\alpha A(z) + \beta B(z)$  is the generating function associated to  $\{\alpha a_n +$ 

2 A(z) · B(z) is the generating function associated to

$$\{c_n\}_{n\geq 0}=\sum^a a_k b_{n-k}$$

Example 2.9. In how many ways can change be given for 30 cents

sing pennies, nickels, dimes, and quarters:

→ Let's look at the generating functions for each currency: Pennies:  $(1+z+z^2+z^3+\cdots)$  Nickels:  $(1+z^5+z^{10}+z^{15}+\cdots)$  Dimes:  $(1+z^{10}+z^{20}+z^{20}+z^{10}+\cdots)$  Quarters:  $(1+z^{25}+z^{20}+z^{75}+\cdots)$  The product of these polynomials is the total number of ways to

 $A(z)B(z)C(z)D(z) = 1 + z + z^{2} + z^{3} + z^{4} + 2z^{5} + \cdots + 18z^{30}$ 

Therefore, there are 18 ways to make change for 30 cents

### 2.9 The Inclusion/Exclusion Principle

Example 2.7. If  $a_n=1$  for all  $n\geq 0$ , then the generating function is This applies to cardinality, area, mass, volume, etc...  $A(z)=1+z+z^2+z^3+\cdots+z^n=\frac{1}{4z^n}=\sum_{n=0}^{\infty}z^n$ How many elements are there in  $A\cup B$  where A and B are finite

 $|A \cup B| = |A| + |B| - |A \cap B|$ 

Now consider three finite sets:  $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$ 

Notation for three finite sets:  $|A_1 \cup A_2 \cup A_3| = \sum |A_i| - \sum |A_i \cap A_j| + |A_1 \cap A_2 \cap A_3|$ 

Theorem 14 (Inclusion/Exclusion), Let  $A_1, A_2, \dots, A_n$  be finite sets.

quadratic cubic polynomial

exponential factorial

4. Average Case Complexity is the average number of operations

used to solve a problem over all inputs of a given size

Theorem 22. Let  $P : \mathbb{R} \to \mathbb{R}$  and  $q : \mathbb{R} \to \mathbb{R}$  be polynomials, then

Consider the problem of making change for n checuts using guarters

dimes, nickels, and pennies using the fewest total number of coins.

The strategy for this problem is defined as the following. At each step.

5.2 Greedy Algorithms

5.2.1 Change Problem

William Farmer

Suject Bhat

Example 2.3. How many ways are there to arrange the letters in 2.5 Counting Integer Solutions The number of different, non-negative integer solutions  $(y_1, y_2, \dots, y_k)$ of the contains.

The number of different, non-negative integer solutions  $(y_1, y_2, \dots, y_k)$ of the contains.

 $TLLHSSEE \rightarrow \frac{8!}{2!2!2!}$ 

Next, determine the possible slots for the "A"'s to go, which are in between each of the letters, as well as at the beginning and end. This leads to a total of

 $\left(\frac{8!}{2!2!2!}\right) \cdot \binom{9}{3}$ 

# 2.3 Binomial Coefficients

Theorem 8 (The Binomial Theorem). Let x and y be variables, and recurrence relation of the form let n be a non-negative integer then

$$(x + y)^n = \sum_{j=0}^{\infty} {n \choose j} x^{n-j} y^j$$

### 2.4 Powersets

The powerset of a set is the set of all its possible subsets.

Example 2.4. How many subsets does the set  $\{1, 2, 3, 4, \dots, n\}$  have? → Let's count sets of size Let  $r_1$  and  $_2$  be the roots of the characteristic equation

•  $1 \Rightarrow \binom{n}{1}$ 

 $\binom{n}{0}+\binom{n}{1}+\binom{n}{2}+\cdots+\binom{n}{n}=(1+1)^n=2^n (\text{Binomial Theorem}) \xrightarrow{\text{$\hookrightarrow$ Assume $a_n=cr^n$. This constraints}} ^{\text{$\hookrightarrow$ Assume $a_n=cr^n$.}}$ 

Example 2.5. Solve  $a_n + a_{n-1} - 6a_{n-2} = 0$ ,  $n \ge 2$   $\hookrightarrow$  Assume  $a_n = cr^n$ . This comes from looking at the simplest possible

2.6 Linear Recursion

Ø. APPM 3170 Suject Bhat

William Farmer  $\left| \bigcup_{i=1}^{n} A_{i} \right| = \sum_{I \subset \{1,2,3,4,...,n\}} (-1)^{|I|+1} \left| \bigcap_{i \in I} A_{i} \right|$ 

### 2.9.1 Derangements

A detargement of  $(1, 2, 3, \cdots, n)$  is any p-non-month that leaves no number in its original position.

For a given set,  $(1, 2, 3, \cdots, n)$ , there are approximately  $\frac{n}{q}$  where q is a tautology. Two statements are logically equivalent if  $p \iff q$  is a tautology.

$$n! \cdot \left(\sum_{i=1}^{n} \frac{(-1)^{i}}{i!}\right)$$

### 3 Logic and Proofs

3.1 Propositional Logic

Before we begin we have to define the syntax of these expressions. Let 3.3.2 Proof by Contraposition the letters  $p,q,r,s,\cdots$  denote the various propositions, while T and F  $p \Rightarrow q \equiv q \lor (\neg p) \equiv \neg p \lor \neg (\neg q)$ . Therefore  $\neg q \Rightarrow \neg p$ , denote the truth value of the statement.

First we define the negation of p, denoted  $\neg p$ . This is expressed as 3.3.3 Proof by Contradiction

The exclusive or of p and q is true when exactly one is true, and 3.3.4 Existence Proofs
also otherwise.

## Table 1: Truth Table for Various Statements 3.2 Propositional Equivalences

A derangements A derangement of  $(1, 2, 3, \dots, n)$  is any permutation of these numbers that is always true is called a tautology, while a statement that is always false is called a contradiction, and a statement that is

 $\binom{m+k-1}{k-1}$ 

 $a_n = c_1a + n - 1 + c_2a_{n-2} + \cdots + c_ka_{n-k} + F(n)$ 

where  $n \ge k$ , F(n) is a function of n only,  $c_i \in \mathbb{R}$ ,  $i = 1, 2, \dots, k$ , and

If F(n) = 0 we call this a homogeneous linear recursion of degree k

Theorem 10. Assume a sequence  $\{a_n\}$  satisfies some degree k linear

 $a_n = c_1 a_{n-1} + c_2 a_{n-2}, n \ge 2$ 

 $r^2 = c_1 r + c_2$ 

If  $r_1 = r_2$ , then  $\exists \{\alpha_1, \alpha_2 \in \mathbb{R} | a_n = (\alpha_1 + \alpha_2 n)r_1^n \}$ 

2 If  $r_1 \neq r_2$  then  $\exists \{\alpha_1, \alpha_2 \in \mathbb{R} | a_n = \alpha_1 r_1^n + \alpha_2 r_2^n \}$ 

Think of this as counting the number of ways to distribute m objects

## 3.3 Methods of Proof

3.3.1 Direct Proof

This style of proof directly proves the statement through application of properties definitions or theorems. It is the most com-

To prove existence we can either choose a constructive approach, or a

expression "p if and only if q." APPM 3170

# he statement, "It is not the case that p." Next we define the conjunction of p and q, denoted $p \wedge q$ . This statement is true when both p and q are true, but like otherwise. The disjunction of p and q is true when either p or q is true, and

also otherwise. It provides the conditional statement is defined by the expression "If p; then q\* non-conditional statement is defined by the expression "If p; then q\* non-conditional statement is similar, except it is defined by the statisfying the conditions, and if it's not constructive, then it has to be non-constructive.

Ø, Suicet Bhat

Discrete Math Notes William Farmer Lemma: If  $n \in \mathbb{Z}, n \geq 0$ , then n cents in change (q, d, n, p), using the while q != 0 fewest coins possible, has at most 2d, 1n, 4p and cannot have 2d+ The amount of change in dnp cannot exceed 24.

5.3 Mergesort

The algorithm is as follows Step One is to split the given list into two equal sublists until each

Step Once is to spit the given list into two equal subsists until each list contains a single element. Step Two is to merge the sublists until they are sorted. 

Stemma: Let  $I_i$ ,  $I_b$  be the two sorted lists of ascerding numbers, where  $I_b$  contains  $n_i$  elements.  $I_i$  and  $I_b$  can be nerged into a single  $I_b$  aprime number can be defined as a positive integer p > 1 if the only look  $I_b$ ,  $I_b$ , using at non  $I_b$ ,  $I_b$  comparisons.

# The worst-case complexity of mergesort is $O(n \cdot \ln(n))$

5.4 Division Algorithm

Every integer can be written as a product of primes uniquely up to the coder of the primes.

For any integers  $a, b \in \mathbb{Z} | a \neq 0, a$  divides b, a|b if  $\exists c \in \mathbb{Z}$  such that If s is a composite integer then n has a prime divisor  $\leq \sqrt{n}$ , and  $\exists c \in \mathbb{Z}$  be positive integers, then there are unique integers a, c, reprime.

Let  $a_b$  be positive integers, used that a=bq+r.  $0 \le c b$  such that a=bq+r. If we consider a fixed b > 1 then  $\exists k \geq 0$  and  $\exists \{(a_0,a_1,\cdots,a_k) \in \{0,1,\cdots,b-1\}\}$  5.6.1 GCD  $[(a_k \neq 0) \land (n = a_k b^k + a_{k-1} b^{k-1} + \cdots + a_0 = \sum_{i=0}^k a_i b^i)]$ 

# 5.4.1 Uniqueness

APPM 3170

5.5 Base b Expansion The following algorithm finds the base b representation of any integer

The complexity of the above algorithm is  $\Theta(loo_n(n))$ 

If a number is not prime, it is composite Every integer can be written as a product of primes uniquely up to

<u>.</u>

For integers  $a, b \in \mathbb{Z}$ , a positive integer c is called the greatest common divisor of a and b if

 (c|a) ∧ (c|b) The representation of any number  $n \in \mathbb{Z}^+ \cup \{0\}$  is unique for each  $2 \cdot (d|a) \wedge (d|b) \Rightarrow (d|c) \Rightarrow (d \leq c)$ Two numbers are relatively prime if their GCD is one.

If a, b, a, r are non-negative integers such that a = ba + r, then the qcd(a, b) = qcd(b, r).If  $a, b \in \mathbb{Z}$  and gcd(a, b) = 1 then  $(\exists \alpha, \beta \in \mathbb{Z}) [1 = \alpha a + \beta b]$ The corollary of the above equation is that if  $a, b \in \mathbb{Z}$ , then  $(\exists \alpha, \beta \in \mathbb{Z}) [\gcd(a, b) = \alpha a + \beta b]$ 

Suject Bhat

case:  $a_{-} = ra_{-}$ ,  $n \ge 1$ ,  $a_{0} = c \rightarrow a_{-} = cr^{n}$ 

$$\hookrightarrow cr^n + cr^{n-1} - 6cr^{n-2} = 0 \rightarrow 1 + r^{-1} - 6r^{-2} = 0$$

 $a_{-} = c_1 2^n + c_2 (-3)^n$ 

Theorem 11. Recall a non homogeneous linear recursion with constant

 $a_n = c_1 a + n - 1 + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n)$ 

 $a_n=c_1a+n-1+c_2a_{n-2}+\cdot\cdot\cdot+c_ka_{n-k}$ 

Any solution to the non-homogeneous linear recursion has the form

and  $b_n$  where  $a_n$  is a particular solution of the non-homogeneous form and  $b_n$  is any solution of the homogeneous form, i.e. the same equation from differential equations with

 $F(n) = (polynomial) \cdot (exponential) = P(n) \cdot S^n$ 

2.6.1 Non-Homogeneous Linear Recursion

with the associated homogeneous form

$$\hookrightarrow r^2 + r - 6 = 0 \rightarrow r_{1,2} = 2, -3$$

So  $a_n=c_12^n$  and  $b_n=c_2(-3)^n$  are solutions. In fact, since they are linearly independent solutions, the general solution is<sup>4</sup> Where q(n) is again a polynomial with degree  $q \le \deg(P)$  is n. 2. When S IS a root of the characteristic equation, then the form is

$$a_n = q(n) \cdot S^n$$
  
gain a polynomial with des  
ot of the characteristic equ  
 $a_n = n^m \cdot q(n) \cdot S^n$ 

1. When S is NOT a root of the characteristic equation of the second

We can also determine these coefficients with  $a_0 = 1$ ,  $a_1 = 2$  giving our Where m is the multiplicity of S as a root of the characteristic quation and q(n) is the san

### Example 2.6. Find the general solution of

form. Then the form is

 $a_n = 3a_{n-1} + 2^n, n \ge 1, a_0 = 1$ 

$$\hookrightarrow$$
 Note that the homogeneous linear recursion form gives us the roots

To find the particular solution, we note that  $F(n) = 2^n$ , which gives

 $h = c2^n$ 

# 2.7 Divide and Conquer Algorithms

The divide and conquer strategy in general is to solve a given problem of size n by breaking the general problem into  $a \ge 1$  sub-problems of

use  $\{a_n\}$  satisfies the non-homogeneous linear recursion when has the form:

Ø.

# 3.3.5 Uniqueness Proofs

First prove  $(\exists x)[P(x) \Rightarrow T]$ Then prove that if  $P(y)\Rightarrow T$  for any y, then show y=x. Else if  $y\neq x$ , show P(y) is false.

3.4 Induction Theorem 15 (The Well-Ordering Principle). Every non-empty

of  $\mathbb{Z}^+$  contains a smallest element.  $\mathbb{Z}^+$  itself is well-ordered. Note,  $\mathbb{Z}^+$ contains no open sets or intervals. **Theorem 16** (The Principle of Mathematical Induction). Let P(n) be 3. Set Complementation: "The complement of  $A^n$  is  $A^c = \{x \in A^n : x \in A$ 

Suppose  $P(1) \Rightarrow T$  and  $\forall k \in \mathbb{Z}^+$  if wherever  $P(k) \Rightarrow P(k+1)$ , then

We can apply DeMorgan's Laws.  $P(n) \Rightarrow T \text{ for all } n \in \mathbb{Z}^+$ ires two steps, the first of which being to prove

# P(1), and the second to prove $P(k) \Rightarrow P(k+1)$

APPM 3170

4 Set Theory

Theorem 17. Definitions 1. A set is a list of elements where repetition and order doesn't 4.2 Set Properties and Functions

3. Two sets are equal if they have exactly the same elements ull set, denoted  $\{\}$  or  $\emptyset$ . Note,  $\{0\}$  is not the empty set 5. A is a subset of B if  $\forall x[x \in A \Rightarrow x \in B]$  is true. We write  $A \subseteq B$ , and  $A \subseteq A$ . Note, A = B if and only if  $A \subseteq B$  and  $B \subseteq A$ .

### 4.1 Operations Between Sets

1. Union: For  $A, B \subseteq u$  we define  $A \cup B = \{x \in u | (x \in A) \lor (x \in B)\}.$ 

2. Intersection: For  $A, B \subseteq u, A \cap B = \{x \in u | (x \in A) \land (x \in B)\}.$ 

2. The difference between A and B is  $A-B=\{x\in u|(x\in A)\land (x\not\in A)\}$ 

6. A is a proper subset of B if A is a subset of B, and  $A \neq B$ . So  $\exists x[x \in B \land x \not\in A]. \ A \subset B$ .

# Discrete Math Notes

5.7 Modular Arithmetic

Fix  $m \ge 2(m \in \mathbb{Z})$  and if  $a, b \in \mathbb{Z}$ , then a is congruent to  $b \mod m, a \equiv b \pmod m$ , if and only if m | (a - b) and  $\exists k \in \mathbb{Z}$  such that  $a - b = b \pmod m$ .  $mk \Rightarrow a = b + mk$ . 1. If  $a \equiv b \pmod{m} \Rightarrow (a = q_1m + r) \land (b = q_2m + r)$ . In other words, an omultiple

and b have the same remainder after dividing by m2. If  $a = b \Rightarrow a \equiv b \pmod{m}$ 3. If  $a \equiv b \pmod{m}$  and  $a, b \in \{0, 1, 2, \dots, m\} \Rightarrow a = b$ . 3. If  $a = \emptyset$ ,  $\bmod m$ ) and  $a, b \in \{\emptyset, 1, 2, \cdots, m\} \Rightarrow a = b$ . 4.  $a \equiv a(mod m)$ 5.  $a \equiv b(mod m) \Rightarrow b \equiv a(mod m)$ 6.  $a \equiv b(mod m) \Rightarrow b \equiv a(mod m) \Rightarrow a \equiv c(mod m)$ 7.  $a \equiv b(mod m) \Rightarrow (a + c) \equiv (b + c)(mod m) \land (ac) \equiv (bc)$ 

# Let $m=11,\mathbb{Z}_{11}=\{x(\mod 11)|x\in\mathbb{Z}\}$ which is equivalent to $\{[0],[1],[2],\cdots,[10]\}$ . Each box is $[x]=\{k\in\mathbb{Z}|k\equiv x(\mod m)\}$ . These are called equivalence rings.

5.7.1 The Space Z...

For every irrational number  $\alpha$ , there are infinitely many rational

6 Graph Theory

٥

 $a_r = 3a_{r-1} \rightarrow r = 3 - a_r = \alpha 3^n \ \alpha \in \mathbb{R}$ 

us that the particular solution has the form

ze  $\frac{n}{b}$  for  $b \ge 1$ . We assume f(n) satisfies  $f(n) = a \cdot f(\frac{n}{b}) + y(n)$ .

We assume f(n) satisfies  $f(n) = a \cdot f\left(\frac{a}{a}\right) + y(n)$ . Let f be an increasing function that satisfies  $f(n) = a \cdot f\left(\frac{a}{a}\right) + c$  where  $a, b, c \in \mathbb{Z}^+$  and  $b \ge 2$ . If  $n|b \Rightarrow \prod_i f(a)$  will be  $O(n^{(np)(a)})$  if  $a > 1\mathbb{Z}$  our time has growth on the order of  $O(\log(n))$ . Furthermore, when a > 1, and  $n = b^i, k = 1, 2, \cdots$  then the time complexity  $f(n) = c_1 \cdot n^{\log(a)} + c_2$  where  $c_i = f(1) + \frac{c_1}{a-1}$  and  $c_2 = -\frac{c_2}{a-1}$ .

William Farmer

 $A_i = \{x \in u | (\exists i \in I)[x \in A_i]\}$ 

 $\bigcap A_i = \{x \in u | (\forall i \in I)[x \subset A_i]\}$ 

 $u|x \notin A$ }.  $u^c = \varnothing$ .  $\varnothing^c$  $\left(\bigcup A_i\right)^c = \bigcap A_i^c$ 

2. If p(z) is a propositional function with domain of speech u (the universe) then  $A = \{ r \in \text{ulp}(x) \}$ , so  $x \in A \Leftrightarrow p(x)$  is true. By definition, the negation of  $x \in A$  is  $x \notin A$ .  $\{(a,b)|(a \in A) \land (b \in B)\}$ 

4 By definition, the only set with no elements is the Empty Set, or A function from A to B is a rule that associates a unique element in B to each element of A, i.e. f: A → B is a function from A to B if (∀a, b ∈ A)[a = b ⇒ f(a) = f(b)]. If f: A → B is a function, then A is called the domain of f, B is

called the commain of f,B is called the domain of f,B is called the codomain, and the range of f is  $f(a) = \{y \in B | (\exists a \in A)|y = f(a)\}$ . Note, by definition, the range is contained in the codomain  $f(a) \subseteq B$ .

### APPM 3170 Ó.

two nodes.

A simple graph is defined as an undirected graph with no loops and io multiple edges.

If a graph is undirected, then the total degree of the vertices is equal

to twice the edges, therefore there must be an even sum of degrees. We also have out degree and in degrees. A graph is called bipartite if it can be written as  $V = V_1 \cup V_2$  where  $V_1 \cap V_2 = \emptyset$ , and every edge is of the form  $\{a,b\} \in G \land a \in V_1 \land b \in V_2$ . A complete bipartite graph has every node in  $V_1$  adjacent to every

We can express graphs as adjacency matrices

5.8 Dirichlet's Approximation Theorem

numbers  $\frac{p}{q}$  such that  $\left|\alpha - \frac{p}{q}\right| < \frac{1}{q^2}$ . Lemma: For any integer  $n \ge 1$  there is a rational number  $\frac{p}{q}$  such that  $\left|\alpha - \frac{p}{q}\right| < \frac{1}{nq}$  where  $1 \le q \le n$ .

A graph can be defined by letting V be a finite, non-empty set of nodes and E be a set of edges. The pair of sets forms a graph. In directed graphs we care about the direction of the nodes, and the order of the pairs in E matter.

Suieet Bhat

William Farmer

Suject Bhat